

MS DryCare

For optimal housing hygiene



MS Schippers

Passion for Farming

Hygiene in livestock housing is essential to Minimize environmental bacteria. Poor hygiene contributes to health problems developing among animals.

A powerful moisture absorbing hygiene powder contributes to optimum health.



Proper housing hygiene helps to ensure good teat condition, udder health and an optimal living environment.



Dry pens contribute to a healthy environment for calves.



A dry pen ensures that piglets get warm quickly.



Scatering the powder in laying nests ensures dry conditions and a healthy climate in the house.

Important aspects of a hygiene powder:

- Moisture absorption – the most important aim of a hygiene powder is to create a dry living environment, so that microorganisms have no chance to develop
- Good distribution – the product must be easy to spread, so that it covers the area well, and does not create dust
- An attractive aroma – plays a role in a pleasant living environment and it can smother or neutralise unpleasant smells
- pH – a hygiene powder must not cause irritation to an animal's teats, skin, hooves or legs



MS DryCare

MS DryCare is a ready-to-use dry powder, suitable for use in animal pens, cubicles, nests etc. to ensure better housing and living conditions. MS DryCare also contains a blend of natural oils based on plant extracts, which have a positive effect on air quality and the animals' airways. The airways of newborn animals open up.

**Very high
moisture
absorption,
>200%**

**Does not
create dust**

**The product is
pH neutral**

Using MS DryCare leads to:

- A dry living environment
- Reduced pathogen pressure
- Improved animal health and comfort


MS DryCare

Cattle



- **Dairy cow cubicles:**
scatter daily on wet patches only

Calves



- **Calving pens and calving areas:**
100 g/m², 2 times per week

Pigs



- **Piglet/sow section:**
50 g/m², 3 times per week
- **Farrowing pens:**
daily, as required

Poultry



- **Broilers:**
100 gr/m², at the start of the round, and then scatter on wet patches
- **Laying nests:**
50 g/m², 1 time every 2 weeks